sPHENIX HCal Simulation Status Overview

Liang Xue

Georgia State University

Minutes from May 1st Simulation Meeting


- 1. Next generation simulation production:
 - Action Item [Chris]: set simulation production in new sPHENIX software environment
 - Action Item [Jin]: Implement Birks Law and more realistic
 Magnetic Field structure in absorber and scintillator for Hcal
 - Action Item [Chris]: Submit production for:
 - 1) different tilt angle as in Liang's slides,
 - 2) pi+ and pi- single particle simulation
 - 3) jet simulation based on pythia jets.
 - QGSP_BERT_HP physics list will be used in this production.
 - Absorber energy will also be saved, so the pion production can also be used to calibrate sampling fraction.
- 2. Action Item [Liang] will analyze the new production data, also with suggested study as in the above action item.

Actual Suggested Study

- What we learn from our first tilted angle scan (2 k events):
 - Mean SF(muon) vs. tilted angle is uniform, less SF(muon) fluctuation for larger tilted angle
 - Energy resolution slightly improved vs. increased tilted angle
- What we suggested for future study:
 - Quantify both high and low side tail of energy spectra for energy leakage/ channeling vs. tilted angle for hadrons
 - Quantify jet performance in term of energy resolution and leakage
 - GEANT4 simulation data production:
 - G4 data for both jets and single particles with 10 k events
 - Detector configuration: Inner HCal 32; Outer HCal 0, 4, 8, 12, 16
 - Detector configuration: Outer HCal 12; Inner HCal 0, 8, 16, 24, 32;
 - Reference: Inner HCal 0, Outer HCal 0

Lower / Higher Side Energy Spectra Tail

- Lower side may due to particle punch-through / channeling
- SF decreases along radius, average SF is ~3%. hadronic showers develops early in outer HCal could be over estimated, and that develops at the far end could be under estimated (Edward).
- Shallow shower has less energy leakage, while deep shower has more energy leakage. This may contribute more to the tails.



- Study energy spectra shape with shower CG cut.
- Correlating low /
 high side tail with
 energy deposited in
 black hole at the
 back of HCal

Next HCal Prototype Simulation

 According John, we are aiming for a beam test in April 2016, and detectors need to be ready for testing at BNL in Jan 2016.

 As long as we have the fixed design of next HCal prototype, we will start to do related simulation study.